

PAPER • OPEN ACCESS

Extremely low frequency ion cyclotron resonances on the surface boundaries of coherent water domains

To cite this article: A Widom *et al* 2021 *IOP Conf. Ser.: Earth Environ. Sci.* **853** 012024

View the [article online](#) for updates and enhancements.

You may also like

- [Formation of Metal Nanoparticles in Ionic Liquids By Low-Energy Quantum Beam Irradiation Technique](#)
Akihito Imanishi
- [Nanometer-Scale Operando Imaging on Ionic and Electronic Transport through Polycrystalline Ceramic Solid Electrolytes](#)
Chun-Sheng Jiang, Ping Liu, Yejing Li et al.
- [Grain Size Effects in Ga-Doped \$\text{Li}_7\text{La}_3\text{Zr}_2\text{O}_{12}\$ Solid Electrolyte for All Solid-State Lithium Ion Batteries](#)
Haneol Park, Won-Sik Kim and Seong-Hyeon Hong



The Electrochemical Society
Advancing solid state & electrochemical science & technology

242nd ECS Meeting

Oct 9 – 13, 2022 • Atlanta, GA, US

Abstract submission deadline: **April 8, 2022**

Connect. Engage. Champion. Empower. Accelerate.

MOVE SCIENCE FORWARD



Submit your abstract



Extremely low frequency ion cyclotron resonances on the surface boundaries of coherent water domains

A Widom¹, J Swain¹ and V I Valenci^{2*}

¹ Physics Department, Northeastern University, Boston MA, USA

² Department of Integrated and Biophysical Medicine, Federiciana Univesita Popolare Via Libero Grassi 11 87100 Cosenza IT, Italy

*vinvalenzi@gmail.com

Abstract. Pure Water under standard pressure, temperature and dilute solution conditions contain coherent Preparata-Del Giudice domains approximately a tenth of a micron in size. Ions of charge $q = Ze$ can be confined to the surface boundaries of coherent water domains. Solving for the motion of ionic charges confined to surfaces in a uniform magnetic field $B = \text{curl}A$ it is shown that there exist resonances scaled by the ionic cyclotron frequency $\omega_c = (qB/Mc)$. The surface classical ionic motions described by the vector potential lagrangian $L = (M/2) \dot{v}^\mu v_\mu + (q/c) \dot{v}^\mu A_\mu$ are integrable for a spherically symmetric surface. The resulting extremely low frequency (ELF) resonant modes are only weakly damped by fluid viscosity due to the small length scales of the confining surface and these modes are of importance in biophysical processes.

1. Introduction

High frequency ionizing radiation is well known to be dangerous to the normal functional behavior of biological organisms [1]. Lower frequencies from radio frequency microwaves to acoustic frequency circuit oscillations at low intensity are considered to have virtually null biological effects. By low intensity we mean that the electromagnetic energy absorption is not sufficient to raise the condensed matter temperature of biological tissues. Typical of measures of biological tissue temperatures are Magnetic Resonance Imaging machines employing magnetic fields of the order of ten Tesla with microwave frequencies of order ten GHz. In detail, the nuclear magnetic resonant frequency of a proton that is followed by the magnetic imaging machine is $\omega_{\text{NMR}} = (g|e|/2Mc)B = \gamma B$, wherein $\gamma \approx 0.2675222$ per nanosec per Tesla.

The Magnetic Resonance Image (MRI) is in fact a picture due to “hot spots” in the living biological tissue. If there tumors within biological tissue, then in the tumor neighborhood the tissue temperature may be well above the normal temperature of the living body. The tumor neighborhood may appear for example as a shadow in the MRI picture. The proton in a hot region diffuses faster than a proton in a cold region. The spin echo experiments probe hot spots by measuring the proton diffusion rates via spin echo damping and turn digitally a temperature map into a picture using a variety computer algorithms. It is however clear that not all tissue damage is accompanied by hot tissue neighborhoods.

Now let us go down perhaps eight orders of magnitude in magnetic field B and in frequency wherein a non-relativistic ion of charge $q = Ze$ and mass M has acyclotron frequency pseudo-vector:



$$\omega_c = -\frac{Z}{M_c} |e| B \quad (1)$$

Fundamental physical mechanisms of the resonant action of an extremely weak alternating magnetic field 40 nano Tesla at the resonant cyclotron frequency in Eq.(1) of a weak 40 milli Tesla static magnetic field in living systems was discussed theoretically [2–5] by Zhadin and coworkers. Extremely low frequency (ELF) resonant effects were also measured [6–8]. At somewhat higher magnetic fields, ELF cyclotron resonant dynamics was measured by Blackman, Liboff and coworkers in other biological systems [9–12]. Finally, in the “phantom”, i.e. nonbiological system of pure water, ELF cyclotron resonances were observed [13] for the H_3O^+ ion.

2. Magnetic dynamics

For a non-relativistic particle moving in a static magnetic field

$$B(r) = \text{curl} A(r) \quad (2)$$

in free space, the vector potential lagrangian describing the motion is given by

$$L(v, r) = \frac{1}{2} M |v|^2 + \frac{q}{c} v * A(r) \quad (3)$$

The momenta

$$p = \frac{dL}{dv} = Mv + \frac{q}{c} A(r) \quad (4)$$

and force

$$f = \frac{dL}{dr} = \text{grad} L = [v A(r)] \quad (5)$$

obey the lagrangian-newtonian rule that the rate of change of momentum is equal to the force
 $p = f$

$$Mv = \frac{q}{c} [\text{grad}(v \cdot A(r)) - (v \cdot \text{grad}) A(r)] \quad (6)$$

or equivalently the magnetic force equation

$$Mv = \frac{q}{c} (v * B) \quad (7)$$

For a magnetic field uniform in space and time, the axial vector cyclotron frequency can be read from Equation (7); It is:

$$\omega_c = -\frac{qB}{Mc} \quad (8)$$

i.e. $\dot{v} = \omega_c \times v$. The mathematics required to repeat this problem for a charge constrained to move on a surface is straight forward.

2.1. Gaussian Surface Constraint

The Gauss differential geometry of two dimensional surfaces in three dimensions is a well known example of differential geometry. For a given coordinate patch one chooses the equation of the surface to be:

$$r = R(x^1, x^2) \Rightarrow dr = e_\mu dx^\mu \quad (9)$$

wherein the Einstein summation convention is employed. The metric distances on the surface depend on the arclength $ds^2 = |d\mathbf{r}|^2$,

$$ds^2 = g_{\mu\nu} dx^\mu dx^\nu \Rightarrow g_{\mu\nu} = e_\mu \cdot e_\nu \quad (10)$$

For a given point on the surface, the vectors ($\mathbf{e}_1, \mathbf{e}_2$) are a basis for the tangent plane while the dual vectors ($\mathbf{e}^1, \mathbf{e}^2$) are a basis for the cotangent plane, wherein, $\mathbf{e}_\mu \cdot \mathbf{e}^\nu = \delta_\mu^\nu$

$$e_\mu = g_{\mu\nu} e^\nu \quad \text{and} \quad e^\mu = g^{\mu\nu} e_\nu \quad (11)$$

The unit vector normal to the surface at a given point is

$$\mathbf{N} = \frac{\mathbf{e}_1 \times \mathbf{e}_2}{|\mathbf{e}_1 \times \mathbf{e}_2|} = \frac{\mathbf{e}_1 \times \mathbf{e}_2}{\sqrt{g}} \quad (12)$$

$$g = |\mathbf{e}_1 \times \mathbf{e}_2|^2 = g_{11}g_{22} - g_{12}g_{21} = \det[g_{\mu\nu}]$$

The vector area element of the surface is thereby

$$d^2\Sigma = (\mathbf{e}_1 dx^1) \times (\mathbf{e}_2 dx^2) = \mathbf{N} \sqrt{g} dx^1 dx^2 = \mathbf{N} d^2\Sigma \quad (13)$$

Any vector tangent or cotangent to the surface at a point may be written as

$$\mathbf{V} = V^\mu \mathbf{e}_\mu = V_\nu \mathbf{e}^\nu \quad (14)$$

When parallel transporting a vector along the surface one must rotate the basis vectors (say) in the tangent plane:

$$\begin{aligned} d\mathbf{V} &= (dV^\mu) \mathbf{e}_\mu + V^\mu (d\mathbf{e}_\mu) \\ d\mathbf{e}_\mu &= \Gamma_{\mu\nu}^\lambda dx^\nu \mathbf{e}_\lambda \\ d\mathbf{V} &= (DV^\mu) \mathbf{e}_\mu \\ DV^\mu &= dV^\mu + \Gamma^{\mu\nu\lambda} V^\nu dx^\lambda \end{aligned} \quad (15)$$

The small rotation of the tangent plane basis vectors ($\mathbf{e}_1, \mathbf{e}_2$) defines the connection coefficients in the second term on the right hand side of Equation (15).

2.2. Equations of Motion

The constraint of having the non-relativistic charged particle moving on a surface can be described by the vector potential lagrangian Eq.(3) confined to the Gaussian surface [14, 15]

$$L(\mathbf{v}, \mathbf{x}) = \frac{M}{2} g_{\mu\nu}(\mathbf{x}) v^\mu v^\nu + \frac{q}{c} v^\mu A_\mu(\mathbf{x}) \quad (16)$$

wherein:

$$\mathbf{V}^\mu = \mathbf{X}^\mu \quad (17)$$

The canonical momenta and force components in the cotangent plane are respectively:

$$\begin{aligned} p_\mu &= \frac{\partial L}{\partial v^\mu} = M g_{\mu\nu}(\mathbf{x}) v^\nu + \frac{q}{c} A_\mu(\mathbf{x}) \\ \mathcal{F}_\mu &= \frac{\partial L}{\partial x^\mu} \\ \mathcal{F}_\mu &= \frac{M}{2} \partial_\mu g_{\alpha\beta}(\mathbf{x}) (v^\alpha v^\beta) + \frac{q}{c} v^\nu \partial_\mu A_\nu(\mathbf{x}) \end{aligned} \quad (18)$$

The lagrangian equations of motion sets the rate of change of momentum equal to the force

$$p_\mu = \int \mu \Rightarrow a^\mu \frac{Dv^\mu}{dt} = v^\mu + \Gamma^\mu_{\lambda\sigma} v^\lambda v^\sigma \quad (19)$$

wherein:

$$\Gamma^\mu_{\lambda\sigma} = \frac{1}{2} g^{\mu\nu} (\partial_\lambda g_{\nu\sigma} + \partial_\sigma g_{\nu\lambda} - \partial_\nu g_{\lambda\sigma}) \quad (20)$$

and a mass times acceleration equal to the magnetic cyclotron force on the ion

$$B_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu \Rightarrow Ma^\mu = \frac{q}{c} B^{\mu\nu} v_\nu. \quad (21)$$

With a tensor rotational velocity

$$\Omega_{\mu\nu} = -\frac{qB_{\mu\nu}}{Mc} \quad (22)$$

one finds cyclotron rotations

$$\frac{Dv^\mu}{dt} + \Omega^\mu_{\nu}(x)v^\nu = 0 \quad (23)$$

In terms of the magnetic field component $B_\perp = \mathbf{N} \cdot \mathbf{B}$ normal to the confining surface,

$$B_{\mu\nu} = \nabla_\mu B_\nu - \nabla_\nu B_\mu = E_{\mu\nu} B_\perp \quad (24)$$

The equations of motion of the charged ion confined to a surface can also be written in Hamilton-Jacobi form.

2.3. The Hamilton-Jacobi Equation

The energy of the confined ion is purely kinetic

$$E = v^\mu \frac{\partial L}{\partial v^\mu} - L = \frac{1}{2} M g_{\mu\nu} v^\mu v^\nu \quad (25)$$

in virtue of Equation (16). Written in terms of the Hamiltonian

$$E = H(p, x) \quad (26)$$

Hamilton's equations of motion are thereby:

$$\dot{x}^\mu = \frac{\partial H}{\partial p_\mu} \quad \text{and} \quad \dot{p}_\mu = -\frac{\partial H}{\partial x^\mu} \quad (27)$$

For the integrable case of a spherical surface in a uniform magnetic field we shall solve the mechanical problem employing the Hamilton-Jacobi equation for the action $W(x, t)$,

$$-\frac{\partial W}{\partial t} = H(p) = \frac{\partial W}{\partial x^\mu} p_\mu \quad (28)$$

the orbits of the ion on the confining surface can be obtained from the first order differential equations $M \dot{x}^\mu = g^{\mu\nu} \partial_\nu W (q/c) A^\mu$ rather than the second order differential Equations (22) and (23).

2.4. Spherical Ordered Water Domain Surface

For a spherical surface of radius R of confinement for one finds cyclotron rotations

$$ds^2 = R^2 (d\theta^2 + \sin^2 \theta d\varphi^2) \quad (29)$$

wherein the uniform magnetic field $\mathbf{B}$ points from the south pole to the north pole. Thus, with $B = |\mathbf{B}|$,

$$B_\theta = B \sin \theta \quad (30)$$

The lagrangian is thereby

$$L = \frac{1}{2}MR^2\dot{\vartheta}^2 + \sin^2\vartheta\dot{\varphi}^2 + \frac{q}{c}(\dot{\vartheta}A_\vartheta + \dot{\varphi}A_\varphi) \quad (31)$$

We choose a gauge for the uniform magnetic field in three dimensional space $A = (1/2)B \times r$ yielding:

$$A_\vartheta = \mathbf{e}_\vartheta \cdot \mathbf{A} = 0 \text{ and } A_\varphi = \mathbf{e}_\varphi \cdot \mathbf{A} = \frac{BR}{2}\sin^2\vartheta \quad (32)$$

Thus, the lagrangian and hamiltonian for the ion confined to the spherical surface are, respectively,

$$L = \frac{1}{2}MR^2\dot{\vartheta}^2 + \sin^2\vartheta\dot{\varphi}^2 + \frac{qBR^2}{2c}\sin^2\vartheta\dot{\varphi} \quad (33)$$

By symmetry, the angular momentum about the magnetic is conserved from rotational symmetry. From the viewpoint of Hamilton Jacobi theory for the action $W(\vartheta, \varphi, t)$, i.e.

$$-\frac{\partial W}{\partial t} = H \quad p_\vartheta = \frac{\partial W}{\partial \dot{\vartheta}}, \quad p_\varphi = \frac{\partial W}{\partial \dot{\varphi}}, \quad \vartheta \quad (34)$$

we show integrability by seeking a solution wherein the energy E and angular momentum J are uniform in time

$$W(\vartheta, \varphi, t) = -Et + J\varphi + S(\vartheta, E, J) \quad (35)$$

Employing the effective “potential” defined by

$$U(\vartheta, J) = \frac{J^2}{2MR^2\sin^2\vartheta} - \frac{1}{2\omega_c J} + \frac{1}{8MR}\omega_c^2\sin^2\vartheta \quad (36)$$

One finds for the action in the energy representation obeying the Hamilton Jacobi equation

$$E = \frac{1}{2MR^2}\partial_{\dot{\vartheta}}^2 S + U(\vartheta, J) \quad (37)$$

The orbit of the ion as a function of time $\vartheta(t)$, $\varphi(t)$ is computed from the action via the implicit equations:

$$\begin{aligned} t &= \frac{\partial S(\vartheta, E, J)}{\partial E} \\ \varphi &= \frac{\partial S(\vartheta, E, J)}{\partial J} \end{aligned} \quad (38)$$

The solution to the Hamilton-Jacobi equation may then be reduced to quadratures

$$S(\vartheta, E, J) = \int_{\vartheta_0}^{\vartheta} q \sqrt{2MR^2E - U(\vartheta, J)} d\vartheta \quad (39)$$

The action corresponding to a closed curve on the spherical surface

$$\tilde{S}(E, J) = \oint q \sqrt{2MR^2E - U(\vartheta, J)} d\vartheta \quad (40)$$

3. Conclusion

This work needs more development in fundamental physics, and opens to a new understanding of many strange biological phenomena, in particular not linear phenomena that influence and example mind-brain-body connections, effects of transdermal drug, that are without chemical reactions as also very low doses of drug far from concentration in blood in the Benveniste Experiment.

Bioelectrical function seems to play a key role not only in heart with ECG but also in all internal organs that seems governed by electrical state of meridians connected that change their electric parameters in answer to any kind of strong and very low signals as showed in the article and in the review by Scalia Valenzi et coll. <http://ibb.kpi.ua/article/view/140255>.

4. References

- [1] Livio Giuliani and Morando Soffritti 2010 Non thermal effects and mechanisms of interactions between electromagnetic fields and living matter (Bologna, Italy : Ramazzi institute) *Eur. J. Oncol.* **5**
- [2] Zhadin M N 1996 Effect of magnetic fields on the motion of an ion in a macromolecule: Theoretical analysis *Biophysics* **4(41)**
- [3] Zhadin M N 1998 Combined action of static and alternating magnetic fields on ion motion in a macromolecule Theoretical aspects *Bioelectromagnetics* **19** 279
- [4] Zhadin M N and Barnes F S 2005 Frequency and amplitude windows at combined action of DC and low frequency AC magnetic fields on ion thermal motion in a macromolecule: Theoretical analysis *Bioelectromagnetics* **26** 323
- [5] Del Giudice E M, Fleischmann, Preparata G and et al. 2005 On the 'unreasonable' effects of ELF Magnetic Fields upon a System of Ions *Bioelectromagnetics* **23** 522
- [6] Novikov V V and Zhadin M N 1994 Combined action of weak constant and variable low-frequency magnetic fields on ionic currents in aqueous solutions of amino acids *Biophysics* **39(1)**
- [7] Zhadin M N, Novikov V V, Barnes F S and et al. 1998 Combined action of static and alternating magnetic fields on ionic current in aqueous glutamic acid solution *Bioelectromagnetics* **19(41)**
- [8] Zhadin M N and Giuliani L 2006 Some problems in modern Bioelectromagnetics *Electromagnetic Biol. Med.* **25** 227
- [9] Liboff A R, Smith S D and McLeod B R 1987 Experimental evidence for 104 cyclotron resonance mediation of membrane transport ed M Blank and E Find *Mechanistic Approaches to Interaction of Electric and Electromagnetic Fields with Living Systems* (New York : Plenum Press) p 109
- [10] Comisso N, Del Giudice E, De Ninno A and et al. 2006 Dynamics of the ion cyclotron resonance effect on amino acids adsorbed at the interfaces *Bioelectromagnetics* **27(16)**
- [11] Liboff A R 1985 Geomagnetic cyclotron resonance in living cells *J. Biol. Phys.* **9(99)**
- [12] Blackman C F, Benane S G, House D E and et al. 1985 Effects of ELF (1-120 Hz) and modulated (50 Hz) RF fields on the efflux of calcium ions from brain tissue in vitro *Bioelectromagnetics* **6(1)**
- [13] D'Emilia E, Ledda M, Foletti A, Lisi A, Giuliani L, Grimaldi S and Liboff A R 2017 Weak Field H_3O^+ ion cyclotron resonance alters water refractive index *Electromagn. Biol. Med.* **36(5562)**
- [14] Landau L D and Lifshitz E M 1987 *Mechanics* Oxford: (Butterworth-Heinemann)
- [15] L.D. Landau and E.M. Lifshitz 2000 *The Classical Theory of Fields* Butterworth-Heinemann Oxford
- [16] Scalia M, Avino P, Sperini M, Viccaro V, Pisani A and Valenzi V I Some Observations on the Role of Water States for Biological and Therapeutic Effects <http://ibb.kpi.ua/article/view/140255>